

European Heart Journal

Risk of malignancy long after acute coronary syndrome in selected urban and rural areas and comparison with smoking risk: The ABC-7* Study on Heart Disease.

--Manuscript Draft--

Manuscript Number:	EURHEARTJ-D-20-04068
Full Title:	Risk of malignancy long after acute coronary syndrome in selected urban and rural areas and comparison with smoking risk: The ABC-7* Study on Heart Disease.
Article Type:	Clinical Research
Keywords:	Acute coronary syndrome; Cancer incidence; Malignancy risk; Urban-Rural; Geographic areas, Survival analysis; UNESCO World Heritage
Corresponding Author:	Giuseppe Berton, MD Conegliano General Hospital Conegliano, TV ITALY
Order of Authors (with Contributor Roles):	Giuseppe Berton, MD Heba Talat Mahmoud Rosa Palmieri Fiorella Cavuto Rocco Cordiano Elisabetta Lorenzon Francesco Bagato
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Conegliano General Hospital
Corresponding Author's Secondary Institution:	
First Author:	Giuseppe Berton, MD
Order of Authors Secondary Information:	
Abstract:	<p>Aims</p> <p>Increased cancer risk has been reported in patients with acute coronary syndrome (ACS). We investigated geographic differences in this risk.</p> <p>Methods and Results</p> <p>We enrolled 586 ACS patients admitted to hospitals in three provinces in the Veneto region of Italy in this prospective study. Patient's residency were classified into three urban and three rural areas.</p> <p>All (except for 3) patients completed the follow-up (22 years or death) and 54% were living in rural areas. Sixteen patients had pre-existing malignancy, and 106 developed the disease during follow-up. Cancer prevalence was 17% and 24% (p=0.05) and incidence of malignancy 16 and 21/1000 person-years for urban and rural areas, respectively. In unadjusted logistic regression analysis, cancer risk increased from urban to rural areas (odds ratio [OR] 3.4;95% confidence interval [CI] 1.7-7.1; p=0.001), with little change from north to south provinces (OR 1.5;95% CI 1.0-2.2; p=0.06). Yet, we found a strong positive interaction between urban-rural areas and provinces (OR 2.1;95% CI 1.2-3.5; p=0.003). These results kept true in the fully adjusted model. Unadjusted Cox regression analysis revealed increasing hazards ratios (HRs) for malignancy onset from urban to rural areas (HR 3.0;95% CI 1.5-6.2; p=0.02), but not among provinces (HR 1.3;95% CI 1.0-2.0; p=0.14). Also, we found a strong positive interaction between geographic areas (HR 2.1;95% CI 1.3-3.5;</p>

	<p>p=0.002), even with a fully adjusted model.</p> <p>Conclusion</p> <p>The results in unselected real-world patients demonstrate a significant geographic difference in malignancy risk in ACS patients, with the highest risk in the north-rural area.</p>
Suggested Reviewers:	
Additional Information:	
Question	Response
Total Word Count:	4330
Word Count Manuscript-only (excluding references):	3999
Did you cite ESC guidelines where appropriate?	yes
As Corresponding Author, I take full responsibility for all information declared in this notification.	Yes
As Corresponding Author, I agree to be the principal correspondent with the Editorial Office, review the edited manuscript and proof, and make decisions about releasing manuscript information to the media, federal agencies, etc.	Yes
All persons who have made substantial contributions to the manuscript (e.g. data acquisition, analysis, or writing / editing assistance), but who do not fulfill authorship criteria, are named with their specific contributions in the Acknowledgements Section of the manuscript.	Yes
All persons named in the Acknowledgements Section have provided the Corresponding Author with written permission to be named in the manuscript.	Yes
If an Acknowledgements Section is not included in the paper then no other persons have made substantial contributions to this manuscript.	Yes
Please enter the names of the authors who did anything else on the manuscript other than what we have listed:	Giuseppe Berton, Heba T.Mahmoud, Rosa Palmieri, Fiorella Cavuto, Rocco Cordiano, Elisabetta Lorenzon, Francesco Bagato.
This manuscript represents valid and substantiated work.	Yes

If asked, I will provide or fully cooperate in obtaining and providing the original data on which the manuscript is based so the editors or their designates can examine it.	Yes
The paper under question is official ESC output being submitted by an ESC Association, Working Group or Council.	No
Each person listed as co-author has been entered as contributing to at least one part of the manuscript	Yes
First Author Secondary Information:	



UNIVERSITÀ DEGLI STUDI DI PADOVA
Dipartimento di Medicina Clinica e Sperimentale
Clinica Medica 4
Policlinico Universitario - Via Giustiniani, 2 - 35128 PADOVA

Professor Filippo Crea
Editor-In-Chief
European Heart Journal.

Padova, September 7, 2020

Dear Prof. Crea,

Please find enclosed our manuscript entitled, "*Risk of malignancy long after acute coronary syndrome in selected urban and rural areas and comparison with smoking risk: The ABC-7* Study on Heart Disease.*", that we are submitting for consideration for publication in the European Heart Journal in the "*clinical research*" section.

We think that this research could be of interest to journal readers, as it shows for the first time a significant geographic difference in malignancy risk in an unselected patients after coronary syndrome during a very long follow-up.

Dr. G. Berton and Dr. F. Cavuto designed the study. Dr. R. Cordiano, Dr. R. Palmieri, and Dr. F. Bagato contributed to the original data collection. Dr. F. Cavuto and Dr. R. Cordiano contributed to data handling and patient follow-up. Dr. G. Berton and Dr. H.T. Mahmoud contributed to the data analysis and interpretation, tables, figures, and manuscript preparation. E. Lorenzon contributed to administrative assistance and figure revision. All authors contributed to ensuring the accuracy of the data analysis.

This paper is not under consideration elsewhere and none of the paper's content has been published previously. All authors have read and approved the manuscript. None of the authors have a financial interest in the subject matter or materials discussed in the manuscript.

I look forward to receiving your comments.

Sincerely,

Giuseppe Berton

ABSTRACT

Aims

Increased cancer risk has been reported in patients with acute coronary syndrome (ACS). We investigated geographic differences in this risk.

Methods and Results

We enrolled 586 ACS patients admitted to hospitals in three provinces in the Veneto region of Italy in this prospective study. Patient's residency were classified into three urban and three rural areas.

All (except for 3) patients completed the follow-up (22 years or death) and 54% were living in rural areas. Sixteen patients had pre-existing malignancy, and 106 developed the disease during follow-up. Cancer prevalence was 17% and 24% ($p=0.05$) and incidence of malignancy 16 and 21/1000 person-years for urban and rural areas, respectively. In unadjusted logistic regression analysis, cancer risk increased from urban to rural areas (odds ratio [OR] 3.4;95% confidence interval [CI] 1.7-7.1; $p=0.001$), with little change from north to south provinces (OR 1.5;95% CI 1.0-2.2; $p=0.06$). Yet, we found a strong positive interaction between urban-rural areas and provinces (OR 2.1;95% CI 1.2-3.5; $p=0.003$). These results kept true in the fully adjusted model. Unadjusted Cox regression analysis revealed increasing hazards ratios (HRs) for malignancy onset from urban to rural areas (HR 3.0;95% CI 1.5-6.2; $p=0.02$), but not among provinces (HR 1.3;95% CI 1.0-2.0; $p=0.14$). Also, we found a strong positive interaction between geographic areas (HR 2.1;95% CI 1.3-3.5; $p=0.002$), even with a fully adjusted model.

Conclusion

The results in unselected real-world patients demonstrate a significant geographic difference in malignancy risk in ACS patients, with the highest risk in the north-rural area.

1
2
3
4
5
6
7
8
9
10
11
12
13

Risk of malignancy long after acute coronary syndrome in selected urban and rural areas and comparison with smoking risk: The ABC-7* Study on Heart Disease.

(*ABC is an acronym for Adria, Bassano, Conegliano, and Padova Hospitals)

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Giuseppe Berton^{1,2}, Heba T.Mahmoud², Rosa Palmieri^{2,3}, Fiorella Cavuto^{2,4}, Rocco Cordiano^{2,3},
Elisabetta Lorenzon², Francesco Bagato^{2,5}.

31
32
33

Authors Affiliations:

1. Department of Cardiology, Conegliano General Hospital, Conegliano, Italy.
2. The ABC Heart Disease Foundation-ONLUS, Conegliano, Italy.
3. Department of Internal Medicine and Cardiology, Adria General Hospital, Adria, Italy.
4. Department of Cardiology, Bassano del Grappa General Hospital, Bassano del Grappa, Italy
5. Department of Cardiology, Feltre General Hospital, Feltre, Italy.

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Correspondence to:

Dr. Giuseppe Berton, MD, FESC.

Cardiology Department, Conegliano General Hospital.

Via Brigata Bisagno, 31015 Conegliano (TV), Italy.

Tel: +390438663613; Fax: +390438663360.

e-mail: giube.s@alice.it

Declarations of interest: none

1
2
3
4
5
6
7
8
9
10
11

Funding sources:

This work was supported by a grant from the Veneto region in Italy (Veneto Region Act no. 748, Venice, May 14, 2015, grant number 2987929). University of Padova (Padova, Italy) contributed to and supported data collection, management, and analysis. The ABC Study on Heart Disease Foundation-ONLUS provided intellectual support for the present study.

12
13

Acknowledgements

14
15
16
17
18
19
20
21
22
23
24
25
26

The authors thank Paola Michelazzo, RN; Jessica Civiero, RN; and nurses from the emergency care units of Conegliano, Adria, and Bassano General Hospitals for their assistance with patient management. We thank Mario Baggio, RN; Daniela Donadel, RN; and Raffaella Frare, RN, for their help handling the data. We thank Renzo De Toni, Ph.D.; Patrizio Buttazzi, Ph.D.; and the general laboratory personnel for assistance collecting laboratory data.

27
28

Authorship and contributors

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Dr. G. Berton and Dr. F. Cavuto designed the study. Dr. R. Cordiano, Dr. R. Palmieri, and Dr. F. Bagato contributed to the original data collection. Dr. F. Cavuto and Dr. R. Cordiano contributed to data handling and patient follow-up. Dr. G. Berton and Dr. H.T. Mahmoud contributed to the data analysis and interpretation, tables, figures, and manuscript preparation. E. Lorenzon contributed to administrative assistance and figure revision. All authors contributed to ensuring the accuracy of the data analysis.

ABSTRACT

Aims

Increased cancer risk has been reported in patients with acute coronary syndrome (ACS). We investigated geographic differences in this risk.

Methods and Results

We enrolled 586 ACS patients admitted to hospitals in three provinces in the Veneto region of Italy in this prospective study. Patient's residency were classified into three urban and three rural areas.

All (except for 3) patients completed the follow-up (22 years or death) and 54% were living in rural areas. Sixteen patients had pre-existing malignancy, and 106 developed the disease during follow-up. Cancer prevalence was 17% and 24% ($p=0.05$) and incidence of malignancy 16 and 21/1000 person-years for urban and rural areas, respectively. In unadjusted logistic regression analysis, cancer risk increased from urban to rural areas (odds ratio [OR] 3.4;95% confidence interval [CI] 1.7-7.1; $p=0.001$), with little change from north to south provinces (OR 1.5;95% CI 1.0-2.2; $p=0.06$). Yet, we found a strong positive interaction between urban-rural areas and provinces (OR 2.1;95% CI 1.2-3.5; $p=0.003$). These results kept true in the fully adjusted model. Unadjusted Cox regression analysis revealed increasing hazards ratios (HRs) for malignancy onset from urban to rural areas (HR 3.0;95% CI 1.5-6.2; $p=0.02$), but not among provinces (HR 1.3;95% CI 1.0-2.0; $p=0.14$). Also, we found a strong positive interaction between geographic areas (HR 2.1;95% CI 1.3-3.5; $p=0.002$), even with a fully adjusted model.

Conclusion

The results in unselected real-world patients demonstrate a significant geographic difference in malignancy risk in ACS patients, with the highest risk in the north-rural area.

Keywords

Acute coronary syndrome; Cancer incidence; Malignancy risk; Urban-Rural; Geographic areas,
Survival analysis; UNESCO World Heritage.

DO NOT
DISTRIBUTE

INTRODUCTION

Cardiovascular disease (CVD) and cancer are the two major causes of mortality worldwide ^{1,2}.

Recently, an increased risk of cancer development has been reported in patients affected by CVD, particularly acute coronary syndrome (ACS) ³⁻⁶. Variation in cancer incidence has been observed between urban and rural areas for several decades, with higher risk of cancer in urban populations ⁷⁻⁹. However, these differences diminished, and even inverted, over the past few years ¹⁰⁻¹³.

To the best of our knowledge, these differences have never been reported in specific populations, such as patients with ACS. In the present study, we investigated the possible difference in malignancy risk in six geographic areas of the Veneto region in Italy in an unselected sample of patients discharged alive after an index hospitalization with ACS and followed up for 22 years. As a comparison, we report the risk of neoplasia in smokers and non-smokers in the same sample of patients.

METHODS

Patients

The ABC Study on Heart Disease is an ongoing prospective investigation designed to represent, as closely as possible, an unbiased population of patients with ACS (www.abcheartdiseasestudy.org/en/). The cohort includes Caucasian patients with definitive ACS, including ST-elevation myocardial infarction (STEMI), non-ST elevation myocardial infarction (NSTEMI), or unstable angina, who were admitted to the intensive care units of three general hospitals in the Veneto region of Italy between June 1995 and January 1998. The original aim of the ABC Study was to monitor these patients with regards to long-term natural history, both non-fatal and fatal events, and causes of death. Another study aim was to investigate the prognostic value of

1 multiple baseline clinical variables. Criteria for diagnosis of ACS included clinical presentation,
2 electrocardiogram findings, and the presence of serum biochemical markers of necrosis ^{14,15}.

3
4 A total of 741 patients were considered eligible upon admission, but 84 patients were excluded
5
6 because they had diseases other than ACS, 23 were excluded due to a lack of baseline data, and 48
7
8 patients were not included in the study because they lived outside the Veneto region. Among the
9
10 586 patients with ACS who were enrolled in the present study, 45 died during the index
11
12 hospitalization. Therefore, the post-discharge follow-up study included 541 patients (Figure 1).
13
14 Each patient received an anonymous code, and no personal data or identifiers were included in the
15
16 baseline or follow-up database.
17
18
19
20

21 **Veneto region and urban-rural classification**

22
23 Veneto is a north-eastern Italian region extending from the Alps to the Adriatic Sea (Figure 2). The
24
25 region is divided into seven provinces, and Venice is the regional capital. The population is
26
27 approximately 5 million, ranking 5th in Italy. The region covers 18 399 km², 6% of the total Italian
28
29 territory, and going south to north it turns from flat to hilly and mountainous. Ninety-five percent of
30
31 the region is rural according to the classification proposed in the national strategic Rural
32
33 Development Plan (Programma di Sviluppo Rurale [PSR] 2014-2020). The present study included
34
35 patients from hospitals in Conegliano-Vittorio Veneto, Bassano, and Adria-Cavarzere, cities located
36
37 in the provinces of Treviso (north), Vicenza (middle), and Rovigo (south), respectively. Urban and
38
39 rural areas were designated for each province, and a residence code that follows the PSR 2014-2020
40
41 coding was assigned to each patient based on their residential address ¹⁶. Thus, areas of residency
42
43 were classified into three urban and three nearby rural areas (Figure 2). The total population of
44
45 these six areas is 586 976 inhabitants, with 24% living in urban areas and 76% living in rural areas
46
47
48
49
50
51
52
53 ^{16,17}.

1
2 In July 2019, at the 43rd session of the UNESCO World Heritage Committee, "Prosecco Hills of
3 Conegliano and Valdobbiadene" was added to the UNESCO World Heritage list as a cultural
4 landscape ¹⁸. This area is in Treviso province, and most of the UNESCO areas overlap the
5
6 geographic area of the ABC Study.
7

8 9 **Measurements and follow-up**

10
11 The study protocol followed the Declaration of Helsinki and was approved by the hospitals' ethics
12
13 committees. All enrolled patients provided written informed consent.
14

15
16 At enrolment, a thorough patient history was collected from medical records and patient interviews.
17

18
19 All of the analysed baseline clinical and laboratory data were obtained during the first 7 days of
20
21 hospitalization in the intensive coronary care unit. The ACS diagnosis criteria were the presence of
22
23 at least two of the following: central chest pain lasting >30 minutes; typical changes in serum
24
25 enzymes, including total creatine kinase (CK) and creatine kinase MB (CK-MB); and typical
26
27 electrocardiogram changes with pathological Q waves and/or localized ST-T changes in at least two
28
29 contiguous leads ¹⁹. Details regarding the measured variables were published previously ^{14,15}.
30
31

32
33 Each patient underwent a clinical check-up 1, 3, 5, 7, 10, 12, 15, 17, 20, and 22 years after
34
35 recruitment. At each recruitment hospital, two cardiologists were responsible for monitoring the
36
37 cohort of patients throughout the follow-up. Data were obtained from scheduled examinations,
38
39 public administrations, hospital records, family doctors, post-mortem examinations, and death
40
41 certificates. The medications received during the index hospitalization and follow-up treatments
42
43 were also recorded. For the present study, the presence of neoplastic disease at the index admission
44
45 and the incidence of new malignancy (i.e., the first documented clinical diagnosis of the disease)
46
47 were recorded. All data after enrolment were recorded prospectively following the protocol of the
48
49 ABC Study on Heart Disease ¹⁴. Baseline data and follow-up data were recorded in two different
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

datasheets. For the present analysis, the datasheets were merged after the completion of 22 years of follow-up.

Statistical analysis

The accrued variables were analysed as continuous variables or proportions. We applied log transformations to correct for positively skewed distributions as appropriate. Measured variables were analysed using the unpaired Student's t-test, and categorical variables using Pearson's chi-squared test. If a patient dropped out before 22 years of follow-up, their data were censored at that time.

We constructed survival curves using cumulative hazards²⁰. As a first step, the homogeneity of malignancy risks in the six geographic areas and smokers was tested using the Breslow-Day test of homogeneity, with $p < 0.05$ indicating dis-homogeneity of the odds ratios (ORs). To study the effect modification, a formal interaction term was assessed for overall malignancy risk between the six geographic areas in unadjusted and fully adjusted logistic regressions. Finally, we used Cox survival regressions including a formal interaction between all areas, and adjusting for the main clinical variables, to estimate the risk of neoplasia onset. Schoenfeld residuals were used to test the proportionality assumption with 95% confidence intervals (CIs). We quantified the risk estimate as the hazard ratio (HR). To graphically show the probability of malignancy and the urban-rural HR for neoplasia onset across the north, middle, and south provinces, we utilized the marginal analysis. We assessed the risks of malignancy prevalence and neoplasia onset in smokers using the same steps for comparison.

Baseline characteristics were summarized as medians and interquartile ranges for continuous variables and numbers and percentages for categorical variables. Unless otherwise indicated, two-

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

tailed P-values <0.05 were considered significant. All statistical analyses were performed using STATA 14 (College Station, Texas, USA).

RESULTS

Study population and baseline characteristics

Of the 586 enrolled patients, 54% were living in rural areas of the three provinces. Table 1 summarizes the main clinical characteristics by area of residency. Both groups shared most of the clinical characteristics (Table 1). All enrolled patients completed the follow-up unless pre-empted by death, except three patients for whom survival was censored before 22 years (two withdrew consent and one moved overseas). At the time of enrolment, 16 patients had previously diagnosed malignancy; one of them died during hospitalization. A total of 106 patients developed the disease during follow-up (Figure 1). The most frequent sites of malignancy were lung (22%), colorectal (19%), prostate (15%), breast (5%), pancreas (5%), and leukaemia (5%).

Overall malignancy prevalence by geographic area

By the end of follow-up, the overall prevalence of neoplasia was 21%. The prevalence was 17% and 24% for the urban and rural areas, respectively ($p=0.05$; Figure 2). The Breslow-Day test of homogeneity demonstrated a significant difference in malignancy prevalence between urban and rural areas across the three provinces (north: OR 3.4, middle: OR 1.6, and south: OR 0.7; $p=0.01$). However, a significant difference was also observed going from south to north in each geographic area (urban: OR 0.5, rural: OR 1.9; $p=0.005$). We found no difference across smoking habit (urban: OR 2.3, rural: OR 2.6; $p=0.80$ and north: OR 2.0, middle: OR 5.8, south: OR 1.9; $p=0.21$).

In the unadjusted logistic regression analysis (Table 2), we observed that the OR for malignancy increases from urban to rural areas, whereas only a slight change in risk was observed from north to

1 south provinces. When a formal interaction term was considered (Table 2), we observed a strong
2 positive interaction between the six geographic areas of residency; the risk increased from urban to
3 rural areas when going from south to north provinces. The interaction analysis showed that the risk
4 of malignancy was 21% vs. 16% in the south, 16% vs. 25% in the middle, and 11% vs. 30% in the
5 north for urban and rural areas, respectively (Table 2 and Figure 3A). Similar results were obtained
6 using a fully adjusted logistic model (Table 2 and Figure 3B). Table 2 also reports the cancer risk
7 by smoking habit in both logistic regression models.
8
9

10 **Malignancy incidence rate by geographic area**

11 The patients discharged alive without neoplasia (n=526) represented 5698 person-years of follow-
12 up. The incidence rate of new malignancy was lower in urban than rural areas (16 and 21/1000
13 person-years, respectively; Figure 1). The Breslow-Day test of homogeneity revealed significant
14 differences in the risk of malignancy onset between urban and rural areas across the three provinces
15 (north: OR 3.8, middle: OR 1.4, and south: OR 0.8; p=0.02). A significant difference was also
16 observed from south to north in each geographic area (urban: OR 0.6, rural: OR 1.8; p=0.01). We
17 found no difference across smoking habit (urban: OR 2.3, rural: OR 2.5; p=0.85 and north: OR 2.0,
18 middle: OR 5.1, south: OR 1.9; p= 0.31).
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 The Nelson-Aalen cumulative hazards estimates for cancer risk across geographic areas indicated
39 that the cumulative hazard appears to decline from south to north areas for urban inhabitants, and
40 increases from north to south for rural inhabitants, with the highest risk in the north-rural area. The
41 cumulative hazard was also higher among smokers than non-smokers (Figure 4).
42
43
44
45
46
47

48 In the unadjusted Cox regression analysis (Table 2), we observed that the HR for malignancy onset
49 increases from urban to rural areas, whereas no change was observed from north to south. When an
50 interaction term was considered, we observed the same strong positive interaction between
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 geographic areas, as the risk increased from urban to rural areas across the south to north provinces.
2 Cox regression-based marginal analysis showed that the risk of malignancy onset was 7 vs. 29 in
3
4 the south, 9 vs. 20 in the middle, and 25 vs. 16 in the north for urban and rural areas, respectively
5
6 (Figure 3A). Similar results were obtained using a fully adjusted cox regression model (Table 3,
7
8 Figure 3B). Table 2 also reports the HR for malignancy onset after ACS by smoking habit in both
9
10 Cox regression models.
11
12

13 **DISCUSSION**

14
15
16 The main result of the ABC-7* Study on ACS is that urban-rural geographic areas have different
17
18 long-term risk of malignancy, even after controlling the survival models for age, gender, and main
19
20 clinical features. This difference supports the hypothesis that the urban-rural geographic area is a
21
22 strong independent effect modifier of malignancy risk in ACS patients.
23
24

25
26 Living in a rural area positively influences the probability of having cancer for patients in the
27
28 northern province, meaning that those who live in the rural areas of this province have a greater risk
29
30 of incurring neoplasia compared to citizens who live in the urban areas. In contrast, living in a rural
31
32 area negatively affects the probability of having cancer in the southern province.
33
34

35
36 Notably, results from the present study also indicate that cancer risk in the north-rural area is even
37
38 higher than the risk associated with smoking (Figure 4). Tobacco smoking is considered a leading
39
40 preventable cause of cancer and cancer deaths ^{21,22}, and strong evidences indicate it significantly
41
42 increases the risk of several malignancies ²³⁻²⁵. In a recent meta-analysis of 19 population-based
43
44 prospective cohort studies included data for 897 021 European and American adults, current-
45
46 smokers had a significantly higher overall risk of developing and dying from cancer (HR 1.44, 95%
47
48 CI 1.28-1.63 and HR 2.19, 95% CI 1.83-2.63, respectively; $P < 0.05$, $I^2 > 75\%$) compared to never-
49
50 smokers ²⁶, which is in concordance with our results based on smoking habits.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Urban-rural variation in cancer risk has been analyzed for several years to emphasize lifestyle
2 dissimilarities, such as smoking and dietary habits, socioeconomic status, and exposures to other
3 risk factors ²⁷⁻²⁹. Yet, homogeneous exposure to the most common risk factors in both urban and
4 rural areas, particularly in developed Western European countries, has recently been reported to be
5 a consequence of economic development, growing homogenization of lifestyle, and increased
6 relocation opportunities ^{28,30}. Thus, the differences in malignancy risk, as well as exposure to
7 different risk factors, seem to depend on health service accessibility, either diagnostic or
8 therapeutic, and secondary prevention strategies, such as national screening programmes ^{10,28,31}.
9 Several reports have documented an elevated cancer risk in the general population in rural
10 communities compared to urban areas ¹⁰⁻¹³. We also recently documented a higher long-term risk of
11 malignancy in patients with ACS compared to the general population ⁴. In our patients, the
12 incidence of cancer was similar in men and women, and higher in older patients than in younger
13 patients ⁴. To the best of our knowledge, the present study is the first to report on the geographic
14 distribution of malignancy in this specific population, ACS patients, with a very long follow-up and
15 very few dropouts. In agreement with the medical knowledge, we found an association between
16 malignancy risk and other important variables, such as age and smoking, but we also found an
17 inverse association between malignancy risk in both urban and rural areas and serum cholesterol as
18 in previous reports ^{32 33}.

19 The explanation for the higher risk in the north-rural area in our cohort of patients, which resulted in
20 approximately 3-times higher risk than in the north urban area, is not yet clear. The six geographic
21 areas considered in this analysis share the same regional and national health service, and it seems
22 unlikely that differences in health system organization can justify the diverse risk of malignancy
23 reported in the present paper. Although recent reports have suggested that the gradual convergence

1 of environmental and economic factors and the changed lifestyle may have contributed to the
2 observed difference in cancer risk, rural residents tend to be older, have a lower income, and less
3 education than urban residents ^{30,34}. Gandinia et al. recently analysed data from 74,989 individuals
4 aged > 35 years and living in 1442 Italian municipalities, who were recruited in 1999-2000 and
5 followed up until 2008. In agreement with our results, they reported that the majority were living in
6 rural areas with lower education level, and that the events of first hospital admission due to
7 cardiovascular diseases or neoplasia were higher in rural residents ³⁵.

8
9
10
11
12
13
14
15
16
17 Notably, the north-rural area of the present study includes 86% of the population living in the rural
18 areas of the Prosecco Hills of Conegliano and Valdobbiadene (UNESCO cultural landscapes),
19 whereas the north-urban area includes 100% of the urban UNESCO population. On the other hand,
20 these rural UNESCO areas overlap 78% of the population of the north-rural area of the ABC 7*
21 Study. This observation sheds light on the higher risk of malignancy in the UNESCO areas than the
22 other geographic areas in the present study.
23
24
25
26
27
28
29
30

31 One of the biggest strengths of this study is derived from the long duration of follow-up with
32 almost no dropouts. However, this study also has limitations. A major limitation of the ABC 7*
33 Study was that, at the time of patient enrolment, percutaneous coronary angioplasty was not yet
34 used to reopen coronary arteries in patients with STEMI. Thus, whether the results may have been
35 altered by early mechanical reperfusion remains uncertain. Another limitation of the study is that
36 the diagnosis of myocardial infarction did not account for troponin measurement, as it was not in
37 use at that time; therefore, we used the rise and gradual decline of creatine kinase and creatine
38 kinase-MB as biochemical markers of necrosis. Nevertheless, these markers of necrosis are still
39 recommended in the absence of troponin measurement ³⁶. Furthermore, all patients in this study
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 were Caucasians, and we cannot generalize the present findings to other populations and ethnic
2 groups.
3

4 To the best of our knowledge, no previous studies have reported such a contrast in malignancy risk
5 between urban and rural areas for a specific population. This prospective study of unselected real-
6 world patients demonstrated a significant difference in the overall malignancy risk, as well as
7 neoplasia onset in different geographic areas of the Veneto region, with the highest risk in the
8 north-rural area and the lowest risk in the north-urban area, among lifelong ACS patients. The risk
9 seems to be higher than that observed with smoking habits.
10
11
12
13
14
15
16
17
18

19 This study emphasizes the importance of implementing a preventive policy based on territory
20 knowledge. Future studies should investigate the aetiological factors related to the higher cancer
21 risk in north-rural areas.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

DO NOT
DISTRIBUTE

REFERENCES

1. Benjamin EJ, Muntner P, Alonso A, et al. Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart Association. *Circulation*. 2019;139(10):e56-e528.
2. Torre LA, Siegel RL, Ward EM, Jemal A. Global Cancer Incidence and Mortality Rates and Trends--An Update. *Cancer Epidemiol Biomarkers Prev*. 2016;25(1):16-27.
3. Cordero A, Lopez-Palop R, Carrillo P, et al. Prevalence and Postdischarge Incidence of Malignancies in Patients With Acute Coronary Syndrome. *Rev Esp Cardiol (Engl Ed)*. 2018;71(4):267-273.
4. Berton G, Cordiano R, Cavuto F, Bagato F, Segafredo B, Pasquinucci M. Neoplastic disease after acute coronary syndrome: incidence, duration, and features: the ABC-4* Study on Heart Disease. *J Cardiovasc Med (Hagerstown)*. 2018;19(10):546-553.
5. Abreu G, Azevedo P, Arantes C, et al. Prevalence of cancer and its impact on prognosis of acute coronary syndrome population. CPC 2016 – Congresso Português de Cardiologia 2016; 2016; Vilamoura, Portugal.
6. Raposeiras-Roubin S, Abu-Assi E, Munoz-Pousa I, et al. Risk of cancer after an acute coronary syndrome according to the type of P2Y12 inhibitor. *Thromb Res*. 2019;174:51-58.
7. Schouten L, Meijer H, Huvneers J, Kiemeny LJJoE. Urban-rural differences in cancer incidence in the Netherlands, 1989–1991. 1996;25(4):729-736.
8. Dey S, Soliman AS, Hablas A, et al. Urban-rural differences in breast cancer incidence in Egypt (1999-2006). *Breast*. 2010;19(5):417-423.
9. Monroe AC, Ricketts TC, Savitz LA. Cancer in rural versus urban populations: a review. *J Rural Health*. 1992;8(3):212-220.
10. Zahnd WE, James AS, Jenkins WD, et al. Rural–urban differences in cancer incidence and trends in the United States. In: AACR; 2018.
11. Kinney AY, Harrell J, Slattery M, Martin C, Sandler RS. Rural-urban differences in colon cancer risk in blacks and whites: the North Carolina Colon Cancer Study. *J Rural Health*. 2006;22(2):124-130.
12. Meilleur A, Subramanian SV, Plascak JJ, Fisher JL, Paskett ED, Lamont EB. Rural residence and cancer outcomes in the United States: issues and challenges. *Cancer Epidemiol Biomarkers Prev*. 2013;22(10):1657-1667.
13. Chow CJ, Al-Refaie WB, Abraham A, et al. Does patient rurality predict quality colon cancer care? A population based study. 2015;58(4):415.
14. Berton G, Citro T, Palmieri R, Petucco S, De Toni R, Palatini P. Albumin excretion rate increases during acute myocardial infarction and strongly predicts early mortality. *Circulation*. 1997;96(10):3338-3345.
15. Berton G, Cordiano R, Cavuto F, Giacomini G, De Toni R, Palatini P. Predictors of ten-year event-free survival in patients with acute myocardial infarction (from the Adria, Bassano, Conegliano, and Padova Hospitals [ABC] study on myocardial infarction). *Am J Cardiol*. 2012;109(7):966-975.
16. Veneto Rd. Programma di sviluppo rurale del veneto 2014-2020. <https://www.regione.veneto.it/web/agricoltura-e-foreste/sviluppo-rurale-2020>. Accessed 30 dicembre, 2019.
17. http://statistica.regione.veneto.it/banche_dati_societa_popolazione.jsp. Accessed 20 August, 2020.

18. UNESCO. Convention concerning the protection of the world cultural and natural heritage. *World Heritage Committee, Forty-third session 2019*; <https://whc.unesco.org/archive/2019/whc19-43com-18-en.pdf>.
19. Pasternak RC, Braunwald E, Sobel BE. Acute myocardial infarction. In: Braunwald E, ed. *Heart Disease: A Textbook of Cardiovascular Medicine* 5th ed. Philadelphia: WB Saunders; 1997:1198-1191 1207.
20. Austin PC, Lee DS, Fine JP. Introduction to the Analysis of Survival Data in the Presence of Competing Risks. *Circulation*. 2016;133(6):601-609.
21. Gallaway MS, Huang B, Chen Q, et al. Smoking and smoking cessation among persons with tobacco-and non-tobacco-associated cancers. *Journal of community health*. 2019;44(3):552-560.
22. Warren GW, Alberg AJ, Kraft AS, Cummings KM. The 2014 Surgeon General's report: "The health consequences of smoking--50 years of progress": a paradigm shift in cancer care. *Cancer*. 2014;120(13):1914-1916.
23. Botteri E, Borroni E, Sloan EK, et al. Smoking and Colorectal Cancer Risk, Overall and by Molecular Subtypes: A Meta-Analysis. *Am J Gastroenterol*. 2020.
24. Larsson SC, Carter P, Kar S, et al. Smoking, alcohol consumption, and cancer: A mendelian randomisation study in UK Biobank and international genetic consortia participants. *PLoS Med*. 2020;17(7):e1003178.
25. Yeo Y, Shin DW, Han KD, et al. Smoking, alcohol consumption, and the risk of thyroid cancer: a population-based Korean cohort study of 10 million people. *Thyroid*. 2020.
26. Ordonez-Mena JM, Schottker B, Mons U, et al. Quantification of the smoking-associated cancer risk with rate advancement periods: meta-analysis of individual participant data from cohorts of the CHANCES consortium. *BMC Med*. 2016;14:62.
27. Schouten L, Meijer H, Huveneers J, Kiemeneij L. Urban-rural differences in cancer incidence in the Netherlands, 1989–1991. *International Journal of Epidemiology*. 1996;25(4):729-736.
28. Stracci F, Cassetti T, Minelli L, et al. Urban-rural differences in male cancer incidence and mortality in the Umbria region of Italy. *Italian Journal of Public Health*. 2007;4(2).
29. Waller R. The combined effects of smoking and occupational or urban factors in relation to lung cancer. *The Annals of occupational hygiene*. 1972;15(1):67-71.
30. Marcellini F, Giuli C, Gagliardi C, Papa R. Aging in Italy: urban-rural differences. *Arch Gerontol Geriatr*. 2007;44(3):243-260.
31. Parikh- Patel A, Bates JH, Campleman S. Colorectal cancer stage at diagnosis by socioeconomic and urban/rural status in California, 1988–2000. *Cancer*. 2006;107(S5):1189-1195.
32. Kitahara CM, Berrington de Gonzalez A, Freedman ND, et al. Total cholesterol and cancer risk in a large prospective study in Korea. *J Clin Oncol*. 2011;29(12):1592-1598.
33. Berton G, Cordiano R, Cavuto F, Bagato F, Mahmoud HT, Pasquinucci M. Association between plasma lipid levels during acute coronary syndrome and long-term malignancy risk. The ABC-4* study on heart disease. *BMC cardiovascular disorders*. 2019;19(1):119.
34. Alfani G, Di Tullio M. Building Regional Distributions of Wealth for the Republic of Venice and for Veneto. In: *The Lion's Share: Inequality and the Rise of the Fiscal State in Preindustrial Europe*. Cambridge University Press; 2019:181-199.

35. Gandini M, Scarinzi C, Bande S, et al. Long term effect of air pollution on incident hospital admissions: Results from the Italian Longitudinal Study within LIFE MED HISS project. *Environ Int.* 2018;121(Pt 2):1087-1097.
36. Thygesen K, Alpert JS, Jaffe AS, et al. Third universal definition of myocardial infarction. *J Am Coll Cardiol.* 2012;60(16):1581-1598.

DO NOT
DISTRIBUTE

Figure legends:

Figure 1. Flow diagram of the study population and progress during follow-up.

*Per 1000-person years. ACS, acute coronary syndrome.

Figure 2. Map of the Veneto region (red arrow) and the location of the six geographic areas.

The respective malignancy prevalence and incidence rates are shown in each area. RV, Rovigo province; TV, Treviso province; VI, Vicenza province.

Figure 3. Graphical representation of the interaction analysis. The probability of malignancy and the hazard ratios for neoplasia onset in urban-rural areas across three provinces are shown for patients followed 22 years after ACS. (A) Unadjusted models. (B) Fully adjusted models. The risk of neoplasia onset by smoking status is also shown for comparison.

Figure 4.

Nelson-Aalen cumulative hazard estimates for cancer risk over 22 years of follow-up by geographic area (A) and smoking habits (B).

Table 1. Baseline characteristics of patients with acute coronary syndrome according to area of residency.

Variable	Overall sample (n=586)	Urban areas (n=268-46%)	Rural areas (n=318-54%)	P-value
Age, years	67(60-75)	68(59-76)	65(60-75)	0.25
Gender, female	31	30	31	0.75
Education above primary school	24	32	17	<0.0001
Body mass index, kg/m ²	25(23-28)	27(23-27)	25(24-28)	0.05
Smoking habit *	66	66	65	0.75
Alcohol use	75	76	75	0.81
Hypertension	50	52	47	0.23
Diabetes mellitus	24	26	22	0.27
Systolic blood pressure, mmHg	120(110-130)	120(110-130)	120(110-135)	0.50
Diastolic blood pressure, mmHg	80(70-80)	75(70-80)	80(70-80)	0.53
Heart rate, beats/min	72(60-83)	73(62-82)	71(60-83)	0.82
Non-ST elevation ACS	37	36	38	0.54
Killip class >1	37	41	34	0.10
Hb, g/dL	14(12-15)	14(12-15)	14(12-15)	0.83
Blood glucose level, mg/dL	122(101-166)	125(104-175)	119(99-156)	0.20
Serum creatinine level, mg/dL	0.9(0.9-1.1)	0.9(0.8-1.2)	0.9(0.9-1.1)	0.59
CK-MB peak [†] , U/L	103(44-212)	104(47-213)	102(43-207)	0.67
Total cholesterol [†] , mg/dL	206(175-242)	203(171-236)	208(177-243)	0.26

The values are presented as median(interquartile range) or percentages.

ACS, acute coronary syndrome; CK-MB, creatine kinase-MB isoenzyme; Hb, haemoglobin.

* Previous smokers and currently smoking patients.

[†] P-values were calculated using log-transformed data.

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


Table 2. Logistic regression analysis of overall malignancy risk and COX regression analysis of the risk of neoplasia onset over 22 years of follow-up after ACS with the interaction for risks between six geographic areas and by smoking habit.


	Unadjusted model			Fully adjusted model		
Logistic regression analysis (n=586)	OR (95% CI)	P-value		OR (95% CI)	P-value	
Urban-rural areas	3.4 (1.7-7.1)	0.001		3.4 (1.6-7.3)	0.001*	
North-south provinces	1.5 (1.0-2.2)	0.06		1.4 (0.9-2.1)	0.11*	
Interaction (urban/rural areas and south to north provinces)	2.1 (1.2-3.6)	0.003		2.1 (1.2-3.6)	0.003*	
Smoking habit	2.5 (1.6-4.1)	<0.0001		2.4 (1.3-4.6)	0.006**	
Cox regression analysis (n=526)	HR (95% CI)	Z	P-value	HR (95% CI)	Z	P-value
Urban-rural areas	3.0 (1.5-6.2)	3.1	0.002	3.2 (1.6-6.6)	3.2	0.001*
North-south provinces	1.3 (1.0-2.0)	1.5	0.14	1.4 (1.0-2.1)	1.8	0.08*
Interaction (urban/rural areas and south to north provinces)	2.1 (1.3-3.5)	3.1	0.002	2.2 (1.4-3.6)	3.2	0.001*
Smoking habit	1.8 (0.8-4.0)	1.5	0.13	2.9 (1.2-7.21)	2.4	0.01**


OR, odds ratio; CI, confidence interval; HR, hazard ratio.

*Adjusted for age, gender, smoking, alcohol consumption, education level, baseline serum cholesterol, and presence of heart failure at admission.


** Adjusted for age, gender, province, alcohol consumption, education level, baseline serum cholesterol, and presence of heart failure at admission.








(A)



(B)



word count: 4330 words.

DO NOT
DISTRIBUTE

The authors do hereby declare that all illustrations and figures in the manuscript are entirely original and do not require reprint permission.

DO NOT
DISTRIBUTE